arXiv:1512.00387v1 [quant-ph] 1 Dec 2015

An approximation formula for the Bloch-Siegert shift of the Rabi model

K. Rapedius,
Max-Beckmann-Str.35, D-76227 Karlsruhe, Germany
e-mail:kevin.rapedius@gmx.de

December 2, 2015

Abstract

So far the Bloch-Siegert shift of the Rabi model has only been calculated numerically or by means of perturbation theory valid in either the weak or strong driving regime only. Recently Yan, Lü, and Zheng [Phys. Rev. A **91**, 053834 (2015)] showed how to reduce the problem to solving a system of three nonlinear equations. Here, we pursue an alternative approach based on a perturbation expansion extrapolation technique. We are thus able to derive an explicit analytical approximation formula for the Bloch-Siegert shift of the Rabi model which is valid for all parameter regimes from weak to strong driving. Comparison with numerically exact results reveals an excellent agreement over the entire driving-strength range.

The Rabi model, which describes a harmonically driven two level system, is a paradigmatic quantum system which finds various physical applications such as atoms in magnetic fields [1, 2], cavity quantum electrodynamics [3], quantum dynamics and transport [4, 5] and quantum information theory [6] to give a few examples. Its Hamiltonian is given by

$$H(t) = \frac{1}{2}\omega_0\sigma_z + \frac{A}{2}\cos(\omega t)\sigma_x \tag{1}$$

where σ_x, σ_y ans σ_z are the Pauli matrices and scaled units with $\hbar = 1$ are used. One finds that due to the harmonic driving the resonance frequency ω_0 of the undriven ($A = 0$) system is changed to a new frequency $\omega_{\rm res}$. The difference $\delta\omega_{\rm BS} = \omega_{\rm res} - \omega_0$ is referred to as Bloch-Siegert shift [7]. This effect is not captured by the celebrated rotating wave approximation described by the Hamiltonian

$$H(t) = \frac{1}{2}\omega_0\sigma_z + \frac{A}{4}\left(\mathrm{e}^{\mathrm{i}\omega t}\sigma_- + \mathrm{e}^{-\mathrm{i}\omega t}\sigma_+\right) \tag{2}$$

which predicts a resonance frequency $\omega_{\rm res} = \omega_0$. The rotating wave approximation (2) is obtained from the Rabi Hamiltonian (1) by inserting $\cos(\omega t) = (\mathrm{e}^{\mathrm{i}\omega t} + \mathrm{e}^{-\mathrm{i}\omega t})/2$, $\sigma_\pm = (\sigma_x \pm \mathrm{i}\sigma_y)/2$ and neglecting the counterrotating terms [8].

Recently Yan, Lü, and Zheng [9] presented a semi-analytical approximation method for calculating the Bloch-Siegert shift $\delta\omega_{\rm BS}$ over the entire range of driving strengths A by means of solving a system of three nonlinear equations, yielding accurate results, in particular for strong driving since it becomes exact in the limit $A/\omega_0 \to \infty$. In the following we will derive a closed-form analytical approximation for the Bloch-Siegert shift $\delta\omega_{\rm BS}$ which is valid from the weak to the strong driving regime and compare it with the semi-analytical results of Yan, Lü, and Zheng [9] and numerically exact calculations [9].

Our starting point is the approximation

$$\begin{aligned} \omega_{\rm res} = \omega_0 \quad &+ \quad \frac{(A/4)^2}{\omega_0} + \frac{1}{4}\frac{(A/4)^4}{\omega_0^3} - \frac{35}{32}\frac{(A/4)^6}{\omega_0^5} \\ &+ \quad \frac{103}{128}\frac{(A/4)^8}{\omega_0^7} + O(A^{10}) \end{aligned} \tag{3}$$

for the resonance frequency obtained by Sanchez and Brandes [10] by means of matrix perturbation theory. The terms up to order six had already been calculated by Shirley [11] using Salwen's [12] Floquet-theory based perturbation approach. Shirley [11] also showed that the Bloch-Siegert-shift in the extreme strong driving limit $A \gg \omega_0$ is given by

$$\delta\omega_{\rm BS} \approx \frac{A}{2.404826} \tag{4}$$

(see also [9]) where the numerical value is obtained as 1/4 of the first zero of the Bessel function J_0.

Viewed as a perturbation problem the system (1) is non-singular and the strong driving limit $A \to \infty$ is equivalent to $\omega_0 \to 0$. Yet the perturbation expansion (3) shows an unphysical divergence for $\omega_0 \to 0$. In the following we will remedy this behaviour by means of an extrapolation technique. While such an extrapolation can in principle be performed by means of rather generally applicable self-similarity-based methods (see [13, 14, 15] and references therein), the present system permits a simpler and more direct alternative approach that we will apply in the following.

We first consider the second order approximation $\omega_{\rm res} = \omega_0 + \frac{(A/4)^2}{\omega_0} + O(A^4)$ obtained from (3) which is also divergent for $\omega_0 \to 0$. However, we can easily obtain an approximation for the squared frequency $\omega_{\rm res}^2 = \omega_0^2 + \frac{1}{8}A^2 + O(A^4)$ valid to the same order of A which has no unphysical divergence for $\omega_0 \to 0$. This way we obtain the second order extrapolation approximation

$$\begin{aligned} \delta\omega_{\rm BS,2} \quad &= \quad \sqrt{\omega_0^2 + \frac{1}{8}A^2} - \omega_0 \\ &\to \quad \frac{A}{\sqrt{8}} \approx \frac{A}{2.8284} \text{ for } \omega_0 \to 0 \end{aligned} \tag{5}$$

for the Bloch-Siegert shift which has the right qualitative behavior (4) for high amplitudes and the same small amplitude behavior as the second order perturbation result. An improved approximation can be obtained by taking the next

term in the perturbation expansion (3) into account and considering the fourth power $\omega_{\rm res}^4 = \omega_0^4 + \frac{1}{4}\omega_0^2 A^2 + \frac{7}{256}A^4 + O(A^6)$ which leads to the fourth order extrapolation approximation

$$\begin{aligned} \delta\omega_{\rm BS,4} &= \left(\omega_0^4 + \frac{1}{4}\omega_0^2 A^2 + \frac{7}{256}A^4\right)^{1/4} - \omega_0 \\ &\to \frac{A}{\left(\frac{256}{7}\right)^{1/4}} \approx \frac{A}{2.4592} \text{ for } \omega_0 \to 0\,. \end{aligned} \tag{6}$$

In an analogous manner we obtain the extrapolation approximations of orders six

$$\begin{aligned} \delta\omega_{\rm BS,6} &= \left(\omega_0^6 + \frac{3}{8}\omega_0^4 A^2 + \frac{33}{512}\omega_0^2 A^4 + \frac{335}{65536}A^6\right)^{1/6} - \omega_0 \\ &\to \frac{A}{\left(\frac{65536}{335}\right)^{1/6}} \approx \frac{A}{2.4094} \text{ for } \omega_0 \to 0\,. \end{aligned} \tag{7}$$

and eight

$$\begin{aligned} \delta\omega_{\rm BS,8} &= \Big(\omega_0^8 + \frac{1}{2}\omega_0^6 A^2 + \frac{15}{128}\omega_0^4 A^4 + \frac{245}{16384}\omega_0^2 A^6 \\ &+ \frac{943}{1048576}A^8\Big)^{1/8} - \omega_0 \\ &\to \frac{A}{\left(\frac{1048576}{943}\right)^{1/8}} \approx \frac{A}{2.4030} \text{ for } \omega_0 \to 0\,. \end{aligned} \tag{8}$$

We see that with increasing order of approximation the asymptotic behavior gets closer to the exact result (4). Note however that, as in perturbation theory itself or any other asymptotic approximation [16], there is no guarantee that the results keep improving if additional higher order terms are included.

In table 1 we compare the 6th and 8th order extrapolated perturbation theory approximations (Eqs. (7) and (8) respectivelly) with numerically exact calulations [9] and the semi-analytical approximation method of Yan, Lü, and Zheng [9]. Over the whole range of parameters we observe a very good agreement between the numerical results and the approximations of Eqs. (7) and (8). For small and moderate driving amplitudes $A \leq 8.5$ the 8th order extrapolated perturbation theory approximation (8) yields the best results of all considered methods, for higher amplitudes $A \geq 11$ the method of Yan, Lü, and Zheng [9] yields slightly better results.

In summary, by means of an asymptotic extrapolation of a perturbation expansion we have derived a closed-form analytical approximation for the Bloch-Siegert shift of the Rabi model which is valid for weak, strong and intermediate values of the driving amplitude.

The author would like to thank Nina J. Lorenz for helpful comments.

A/ω_0	Numerical [9]	Eq. (7)	Eq. (8)	Approx. in [9]
1.0	0.063224	0.063219	0.063224	0.063268
3.5	0.707959	0.706106	0.708068	0.716200
6.0	1.641809	1.637358	1.642716	1.649924
8.5	2.637787	2.631189	2.639640	2.640075
11.0	3.653740	3.645118	3.656504	3.652351
13.5	4.678502	4.667893	4.682141	4.675271
16.0	5.707919	5.695333	5.712406	5.703825
18.5	6.740093	6.725536	6.745409	6.735637
21.0	7.774035	7.757510	7.780169	7.769474

Table 1: Comparison of relative Bloch-Siegert shifts $\delta\omega_{\rm BS}/\omega_0$ for various driving amplitudes A calculated by different methods: Numerically exact calulations [9], 6th order extrapolated perturbation theory approximation (Eq. (7)), 8th order extrapolated perturbation theory approximation (Eq. (8)) and the semi-analytical approximation method of Yan, Lü, and Zheng [9].

References

[1] I. Rabi, Phys. Rev. **49**, 324 (1936).

[2] I. Rabi, Phys. Rev. **51**, 652 (1937).

[3] C. Cohen-Tanoudji, J. Dupont-Roc, and G. Grynberg, *Photons and Atoms: Introduction to Quantum Electrodynamics* (Wiley, New York, 1997).

[4] T. Dittrich, P. Hänggi, G.-L. Ingold, B. Kramer, G. Schön, and W. Zwerger, *Quantum Transport and Dissipation* (Wiley–VCH, Weinheim, 1998).

[5] L. Henriet, Z. Ristivojevic, P. P. Orth, and K. Le Hur, Phys. Rev. A **90**, 023820 (2014).

[6] P. Kok and B. W. Lovett, *Introduction to optical quantum information processing* (Cambridge University Press, Cambridge, 2010).

[7] F. Bloch and A. Siegert, Phys. Rev. **57**, 522 (1940).

[8] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, 1997).

[9] Y. Yan, Z. Lü, and H. Zheng, Phys. Rev. A **91**, 053834 (2015).

[10] B. N. Sanchez and T. Brandes, Ann. Phys. **10**, 569 (2004).

[11] J. H. Shirley, Phys. Rev. **138**, B979 (1965).

[12] H. Salwen, Phys, Rev. **99**, 1274 (1955).

[13] V. I. Yukalov, E. P. Yukalova, and S. Gluzman, Phys. Rev. A **51**, 96 (1998).

[14] V. I. Yukalov and S.Gluzman, Int. J. Mod. Phys. B **18**, 3027 (2004).

[15] S. GLuzman and V.I. Yukalov, Euro. Jnl of Applied Mathematics **25**, 595 (2014).

[16] M. H. Holmes., *Introduction to Perturbation Methods* (Springer, New York, 2013).